Search for CP Violation in Hyperon Decays: Preliminary Results from the HyperCP Experiment at Fermilab

C. M. Jenkins
Department of Physics
University of South Alabama
Mobile, AL 36688

for the
HyperCP Collaboration

March 19, 1998

This work supported in part by:
DoE grant: DE-FG02-96ER40970

E871 (HyperCP) Collaboration

A. Chan, Y.C. Chen, C. Ho, J.Sheng†, P.-K. Teng,
C. Yu† Z. Yu†
Academia Sinica, Tiawan

W.-S. Choong, G. Gidal, K.-B. Luk*, B. Turko, P. Zyla
University of California and Lawrence Berkeley National Laboratory

M.Crisler, C. James, J. Volk
Fermi National Accelerator Laboratory

J. Felix, G. Lopez, M. Sosa
University of Guanajuato, Mexico

R. A. Burnstein, A.Chakravorty, J. Drapala,
D. M. Kaplan, L.M. Lederman,W. Luebke, D. Rajaram, H. A.
Rubin, C. White, S. White
Illinois Institute of Technology

N. Leros, J.-P. Perroud
University of Lausanne, Switzerland

V. Papavassiliou
New Mexico State University

H. R. Gustafson, M. J. Longo, F. Lopez
University of Michigan

K. Clark, C. M. Jenkins
University of South Alabama

E. C. Dukes*, C. Durandet, M. Huang, K. Nelson
University of Virginia

* co-spokespersons
† Visitors form IHEP, Beijing
Post-Doc
Graduate Student

The HyperCP Experiment took data at the Fermi National Accelerator Laboratory.

- The experiment has took data in the 1997 fixed target run at the Fermilab outside Chicago, IL.
- Results presented here are PRELIMINARY .
- The HyperCP Experiment's primary Physics goal is to look for CP violation in the decays of Ξ hyperons.

$$\begin{array}{lcl} \Xi^- & \to & \Lambda\pi^- \\ \quad\Lambda & \to & \mathrm{p}\ \pi^- \end{array} \qquad \begin{array}{lcl} \overline{\Xi}^+ & \to & \overline{\Lambda}\pi^+ \\ \quad\overline{\Lambda} & \to & \overline{p}\pi^+ \end{array}$$

CP violation has only been observed in the decays of K^o's

- Other Physics to be explored by the HyperCP Experiment include:
  - Search for rare decays:
    - $\mathrm{K}^\pm \to \pi^\pm\mu^+\mu^-$.
    - $\mathrm{K}^- \to \pi^+\mu^-\mu^-$.
    - $\Xi^- \to \mathrm{p}\ \mu^-\mu-$
  - Search for CP violation in $\mathrm{K}^\pm \to \pi^+\pi^-\pi^\pm$ decays.
- All Physics uses a "left-right" trigger for the spectrometer.
  - An event with at least one particle with the beam sign and at least one particle with opposite the beam sign.

CP Violation in the K^o System

- K^o and $\overline{K}^o$ are produced via the strong interaction.
- They decay via the weak interaction.
  - The weak (decay) states are not eigenvectors of the strong (production) states.
  - Results in quantum mechanical mixing.
- Assume that weak interaction is symmetric under CP.
- Expect two CP eigenstates:

$$\begin{array}{ll} |K_1\rangle = \frac{1}{\sqrt{2}}(|K^o\rangle + |\overline{K}^0\rangle) & \text{CP even} \\ |K_2\rangle = \frac{1}{\sqrt{2}}(|K^o\rangle - |\overline{K}^0\rangle) & \text{CP odd} \end{array}$$

- These are CP eigenstates, and decay in to CP eigenstates

$$\begin{array}{llll} CP|K_1\rangle = +1|K_1\rangle & |K_1\rangle \to 2\pi & \text{CP even} \\ CP|K_2\rangle = -1|K_2\rangle & |K_2\rangle \to 3\pi & \text{CP odd} \end{array}$$

- As a result, one expects to see only two types of decays:

Decay	Lifetime	
$k_{short} \to \pi^+\pi^-$	$\tau = 0.892 \times 10^{-10}$ sec	CP even
$k_{long} \to \pi^+\pi^-\pi^o$	$\tau = 5.17 \times 10^{-10}$ sec	CP odd

- In 1964 it was discovered that the long lived $|K_2\rangle$ state could decay into the 2π state!
  - The asymmetry is on the order of 3×10^{-3}.
  - **CP violation has been observed only in the K^o system.**
- CP violation in the K^o system has also been observed in the asymmetry in the decay rates of:

$$K_{long} \to e^+\nu_e\pi^-$$
$$K_{long} \to e^-\overline{\nu_e}\pi^+$$

Physics of CP Violation in Ξ decays

- The non-leptonic decay of the Ξ hyperon is described in terms of the Lee-Yang variables:

$$\alpha = \frac{2Re(S^*P)}{|S|^2 + |P|^2}$$

$$\beta = \frac{2Im(S^*P)}{|S|^2 + |P|^2}$$

$$\gamma = \frac{|S|^2 - |P|^2}{|S|^2 + |P|^2}$$

- Under CP operation, α and β change signs, γ is unchanged.
- The decay distribution for the daughter spin $\frac{1}{2}$ baryon in the rest frame of the parent hyperon is:

$$\frac{dP}{d\Omega} = \frac{1}{4\pi}(1 + \alpha \vec{P}_p \cdot \hat{p}_d)$$

- Where in the rest frame of the parent:
  - $\vec{P}_p$ is the parent hyperon polarization.
  - $\hat{p}_d$ is the daughter baryon momentum direction.
- The daughter baryon is polarized according to:

$$\vec{P}_d = \frac{(\alpha + \vec{P}_p \cdot \hat{p}_d)\hat{p}_d + \beta(\vec{P}_p \times \hat{p}_d) + \gamma(\hat{p}_d \times (\vec{P}_p \times \hat{p}_d))}{(1 + \alpha \vec{P}_p \times \hat{p}_d)}$$

- Where:
  - α, β, γ are the Lee-Yang variables for the parents.

- The HyperCP Experiment uses unpolarized Ξ particles.
  - (Use 0^o production angle for Ξ.)
- As a result the daughter Λ polarization is:

$$\vec{P}_\Lambda = \alpha_\Xi \hat{p}_\Lambda$$

- The polarization of the Λ may be measured by the decay asymmetry:

$$\boxed{\frac{dP}{d\Omega} = \frac{1}{4\pi}(1 + \alpha_\Lambda \vec{P}_\Lambda \cdot \hat{p}_p)}$$

- Where
  - $\hat{p}_p$ is the direction of the proton (anti-proton) in the Λ ($\overline{\Lambda}$) rest frame.
  - Note the polarization of the Λ is : $\vec{P}_\Lambda = \alpha_\Xi \hat{p}_\Lambda$



- The asymmetry in the proton (anti-proton) decay distribution is:
  - (After integrating over the azimuthal angle.)

$$\boxed{\frac{dP}{d(\cos(\theta))} = \frac{1}{2}(1 + \alpha_\Lambda \alpha_\Xi \cos(\theta))}$$

- Note the "Z" axis in the Λ rest frame is defined by the Λ direction in the Ξ rest frame.
- So the "θ" is the polar angle of the proton (or anti-proton) in the Λ rest frame.
- Any differences in the angular distributions of the proton from Ξ^- decays and anti-proton from $\overline{\Xi}^+$ decays is a signature of CP Violation.



- The combined asymmetry parameters for the Ξ and Λ will be determined from the angular distributions.
- A CP asymmetry may be constructed:

$$\mathcal{A} = \frac{\alpha_\Lambda \alpha_\Xi - \alpha_{\overline{\Lambda}} \alpha_{\overline{\Xi}}}{\alpha_\Lambda \alpha_\Xi + \alpha_{\overline{\Lambda}} \alpha_{\overline{\Xi}}}$$

- The HyperCP experiment collected 67×10^9 triggers to tape.
- This data should contain roughly 1.31×10^9 Ξ^- reconstructible decays and 2.84×10^8 $\overline{\Xi}^+$ reconstructible decays.
- This results in a overall sensitivity of 2.0×10^{-4} for the CP asymmetry $\mathcal{A}$.
- Theoretical predictions using the Standard Model predict a value for the CP asymmetry $\mathcal{A}$ of -0.7×10^{-4} to -8.8×10^{-4}.
  - Using other non-standard models will widen this range or even exclude CP violation in Ξ decays.



- HyperCP spectrometer: Plan View.
- 800 GeV/c primary proton beam hits target.
- 150 GeV/c secondaries exit the Hyperon magnet.
  - Negative mode secondary beam contains Ξ^-.
  - Positive mode secondary beam contains $\overline{\Xi}^+$.
- Four PWC's upstream an analysis magnet, Four PWC's downstream.
- Two trigger hodoscopes to form "left-right" trigger.

- Calorimeter used for halo muon rejection.
- Muon identification down stream used for rare decay trigger.



- HyperCP spectrometer: Elevation View.
- The secondaries are bent vertically.



- Event display of a typical Ξ event.
- Tracking is used to draw tracks.
- Hits used by tracking are colored "red".
- Track number and total momentum are calculated.
- Hits in the trigger hodoscope are colored red.
- Calorimeter has percent dynamic range color coded.
- Ξ and Λ mass are calculated.

$$\Xi^- \rightarrow \Lambda\pi^- \qquad\qquad \overline{\Xi}^+ \rightarrow \overline{\Lambda}\pi^+$$
$$\Lambda \rightarrow p\pi^- \qquad\qquad \overline{\Lambda} \rightarrow \overline{p}\pi^+$$



- The invariant mass Ξ^- and $\overline{\Xi}^+$ for 0 mRad production angle.
- These are events that have been processed by the geometric and kinematic fits.
  - Geometric fit: fits events to the one kink and neutral vee topology.
  - Kinematic fit: Assumes the vee is a Λ decay.
    - Conserves momentum in Λ rest frame.
    - Constrains mass to that of Λ.
- Cuts include:
  - $125 < P_\Xi < 250$ GeV , $100 < P_\Lambda < 215$ GeV, $75 < P_{Proton} < 185$ GeV, $P_{\pi_\Xi} < 65$, $P_{\pi_\Lambda} < 55$ GeV
  - $\chi^2_{GFit} < 2$, $\chi^2_{KFit} < 50$
  - -0.6 < Y_{coll} , 0.68 cm, -0.75 < X_{coll} < 0.85 cm, -0.25$\leq X_{tgt} \leq$ 0.32 cm
  - -2075 < V_Ξ < -950 cm , 100 < D_Λ, 1.1573 < M_Λ < 1.1564 GeV

$$\Xi^- \rightarrow \Lambda\pi^- \qquad\qquad \overline{\Xi}^+ \rightarrow \overline{\Lambda}\pi^+$$
$$\Lambda \rightarrow p\pi^- \qquad\qquad \overline{\Lambda} \rightarrow \overline{p}\pi^+$$



- Resolution of the HyperCP spectrometer is very good: 1.39 MeV/c^2 for the $\Xi^-/\overline{\Xi}^+$ mass.



- Monte Carlo of Ξ mass resolution.
- Ξ events are generated and are traced through the hyperon magnets.
- Ξ's are allowed to decay via $\Xi \rightarrow \Lambda\pi \rightarrow p\pi\pi$ mode.
- Events that satisfy the trigger are reconstructed with the same tracking, geometric and kinematic fitting programs as the data.
- This plot contains the invariant mass distribution as calculated from the kinematic fit.
- The resolution is 1.39 MeV/c^2 from the Monte Carlo



- Direction cosines for the proton in the Λ rest frame along the spectrometer axes for Ξ^- vs $\overline{\Xi}^+$.
- These are related to the Ξ polarization.
- The Ξ's were selected according to the previously outline cuts.
- Recall the spectrometer geometry:
  - Z axis is the beam direction.
  - X axis is parallel to the Hyperon magnet magnetic field.
  - Y axis is the polarization direction.
- Figure A is the direction cosine along the X direction.
- Figure B is the direction cosine along the Y direction.
- Figure C is the direction cosine along the Z direction.
  - The difference in the distributions is due to the difference in the production of Ξ^- vs $\overline{\Xi}^+$.

Very Preliminary!



- The cosine of the proton's direction in the Λ rest frame.
- This is the plot studied for CP violation in hyperon decay.
- This plot contains a TINY fraction of the HYPERCP collaboration sample.
- No systematic studies made.
- No corrections for the differences between production of Ξ^- vs $\overline{\Xi}^+$.

$K^{\pm} \rightarrow \pi^{+}\pi^{-}\pi^{\pm}$



- Uses a three prong vertex algorithm.
- The cuts used are:
  - $\chi^2 \leq 10$, 0.400 GeV $< M_K <$ 0.600 GeV, -2200 $<$ Z $<$ -950 cm.
  - $P_{\pi_1} < 0.13$ GeV, $P_{\pi_2} < 0.13$ GeV, $P_{\pi_3} < 0.13$ GeV.
  - $M_{\pi_1\pi_2} < 0.36$ GeV, $M_{\pi_1\pi_3} < 0.36$ GeV, $M_{\pi_2\pi_3} < 0.36$ GeV.
  - $120 < P_K < 250$ GeV.
  - Plus acceptance cut (for pions to go through the analyzing magnet).

$K_{short} \rightarrow \pi^+\pi^-$



- K_{short}'s are produced at the collimator exit.
- Use Same-sign and Opposite-sign track pair that have distance of closest approach.

$$\Omega^- \rightarrow \Lambda\pi^- \qquad\qquad \overline{\Omega}^+ \rightarrow \overline{\Lambda}\pi^+$$
$$\Lambda \rightarrow p\pi^- \qquad\qquad \overline{\Lambda} \rightarrow \overline{p}\pi^+$$



- $\Omega^- \rightarrow \Lambda K^- \rightarrow p\pi^- K^-$.
- This is the same topology as the $\Xi^- \rightarrow \Lambda\pi^- \rightarrow p\pi^-\pi^-$.
- Use the same geometric fit (for the decay geometry).
- And the same kinematic fit (for the Λ decay).
- The Ω^- has a mass of approximately 1.672 GeV/c^2.

Conclusions

- The HyperCP experiment has collected the largest Ξ^- and $\overline{\Xi}^+$ sample in the world.
  - The experiment recently concluded its first round of data collection at Fermilab (September 1997).
  - The experiment collected 67×10^9 triggers.
  - 1.31×10^9 reconstructible Ξ^-
  - 2.84×10^8 reconstructible $\overline{\Xi}^+$
  - The HyperCP experiment anticipates a sensitivity of the CP violating asymmetry $\mathcal{A}$ of 2.0×10^{-4}.
  - The experiment should have:
    * 2.62×10^7 reconstructible Ω^-.
    * 5.68×10^6 reconstructible $\overline{\Omega}^+$.
- The experiment is in the early analysis stage.
- The experiment has processed through tracking about 8% of its data by March 1998.
- The HyperCP experiment has been approved for more beam time in the next fixed target run at Fermilab in 1998.
  - The experiment will increase its data sample by a factor of four.

Experimental Upper Limits on CP Violations in Hyperon Decay

- Only experimental upper limits exist for CP violation in hyperon decays.
- Experiments have looked for CP violation in $\Lambda/\overline{\Lambda}$ decays.

$\overline{p}p \to \Lambda X, \overline{p}p \to \overline{\Lambda} X$	0.02	$\pm$	0.14	R608
$e^+e- \to J/\psi \to \Lambda\overline{\Lambda}$	0.01	$\pm$	0.10	DM2
$\overline{p}p \to \Lambda\overline{\Lambda}$	0.013	$\pm$	0.022	PS185
Recomputed World Average	0.013	$\pm$	0.021	

- The theoretically expected value for the CP asymmetry in the $\Lambda/\overline{\Lambda}$ system is $-0.3 \times 10{-}4$ to -4.0×10^{-4}.
- These experiments are not sensitive enough to see CP violation that is consistent with the Standard Model.

- R608: P. Chauvat et al., Phys. Lett **B163** (1985) 273.
- DM2: M. H. Tixier et al., Phys. Lett. **B212** (1988) 523.
- PS 185: P. D. Barnes et al., Phys. Lett. **B199** (1987) 147.

Λ and Ξ Hyperons

Λ

- Quark content: up, down, strange.
- Electric charge: 0 (neutral)
- Mass: 1.115684 ± 0.000006 GeV/c^2.
- Lifetime: $(2.632 \pm 0.020) \times 10^{-10}$ sec.
- Asymmetry: $\alpha = 0.642 \pm 0.013$.
- Major decay modes:
  - p π^-, Br = 0.639
  - n π^o, Br = 0.358

Particle Data Group, Phys. Rev. (**D54**)51.

Ξ^-

- Quark content: down, strange, strange.
- Electric charge: -1.
- Mass: 1.32132 ± 0.00013 GeV/c^2.
- Lifetime: $(1.639 \pm 0.015) \times 10^{-10}$ sec.
- Asymmetry: $\alpha = -0.456 \pm 0.014$.
- Major decay modes:
  - $\Lambda\pi^-$, Br = 0.99887

Particle Data Group, Phys. Rev. (**D54**)54.